

Bulk Fields in Delocalized Dilatonic p -Branes

Donam Youm¹

ICTP, Strada Costiera 11, 34014 Trieste, Italy

Abstract

We study localization properties of various bulk fields on a dilatonic p -brane which is delocalized along its transverse directions except one. We find that all the bosonic and fermionic bulk fields can be localized on the delocalized dilatonic p -brane in a strict sense, namely the Kaluza-Klein zero modes of the bulk fields are normalizable and are localized around the brane, for any values of the dilaton coupling parameter.

July, 2001

¹E-mail: youmd@ictp.trieste.it

Recently, there has been active interest in brane world scenarios [1, 2, 3, 4], because they provide possible attractive solutions to the hierarchy and the cosmological constant problems. In brane world scenarios, it is generally assumed that only gravity lives in the bulk and all the fields of the Standard Model are confined within the brane worldvolume. Such point of view may be natural in string theories, where matter fields on the brane can be identified as open string modes and the gravity in the bulk is a closed string mode. But when we consider the embeddings of brane world scenarios into supergravity theories, we usually encounter various bulk matter fields, especially those compactified from 10 or 11 dimensions if we are interested in effective supergravity descriptions of string theories. So, it would be of interest to study various bulk matter fields in brane world models. (The previous studies on various bulk fields in the brane world scenarios can be found, for example, in Refs. [5, 6, 7, 8, 9, 10, 11, 12, 13, 14].)

In the case of the Randall-Sundrum (RS) domain wall with noncompact extra space [4], it was shown [6, 7, 8, 9, 10, 11, 12] that it is impossible to localize bulk gauge fields and bulk fermionic fields on the domain wall through purely gravitational mechanism. (Cf. See also Refs. [15, 16, 17].) It was also proposed that a $U(1)$ field on the brane originates rather from two bulk two-form potentials [18, 19] and alternative methods for localizing the bulk $U(1)$ field were also proposed [20, 21]. In our previous works [22, 23, 24], we showed that dilatonic domain walls can also localize bulk gravity, provided the tension of the wall is positive. Unlike the (nondilatonic) RS domain wall, the dilatonic domain walls are shown [23] to localize all the bosonic and fermionic bulk fields, provided the dilaton coupling parameter is large enough in some cases. However, some of bulks fields are not localized on the dilatonic domain walls in a strict sense. Namely, although the Kaluza-Klein (KK) zero modes of all the bulk fields are normalizable, the KK zero modes of some bulk fields are evenly distributed along the extra spatial direction or localized away from the wall. Another problem with the dilatonic domain walls which localize gravity is that there are naked singularities at finite distance away from the wall.

In Ref. [25], we studied localization properties of bulk gravity on dilatonic p -branes in arbitrary spacetime dimensions. We showed that generally bulk gravity cannot be localized on a p -brane (with the codimension higher than or equal to three) unless the transverse space is truncated, for example, through the introduction of another p -brane at finite distance from the existing one. However, when all the transverse directions except one are delocalized, the bulk gravity is shown to be localized on the brane, if a parameter in the p -brane harmonic function is chosen such that there are curvature singularities on both sides of the brane at finite coordinate distance. Such singularities of the delocalized p -branes are less harmful than the naked singularities of the dilatonic domain wall in a sense that massive matter with finite energy can never reach the singularities, being repelled by the singularities. The warp factor, which sets the energy scale on the hypersurface, of such delocalized brane solution increases as one moves away from the brane, asymptotically approaching infinity as the singularities are approached. So, one may regard such delocalized brane as the TeV brane. In this note,

we study the localization properties of various bulk fields on the delocalized dilatonic p -branes. We find that all the bosonic and fermionic bulk fields are localized on the brane in a strict sense, namely not only the KK zero mode is normalizable but also its field distribution is localized around the brane.

We begin by discussing delocalized dilatonic p -brane solution studied in Ref. [25]. Just as in Ref. [25], we regard the p -brane as a solitonic brane magnetically charged under the field strength F_n of the rank $n = D - p - 2$, although such assumption is not essential for our discussion. The action for a dilatonic p -brane in D -dimensional spacetime with an arbitrary dilaton coupling parameter a_p is

$$S_p = \frac{1}{2\kappa_D^2} \int d^D x \sqrt{-G} \left[\mathcal{R} - \frac{4}{D-2} (\partial\phi)^2 - \frac{1}{2 \cdot n!} e^{-2a_p\phi} F_n^2 \right]. \quad (1)$$

The solution for the extreme dilatonic p -brane with the longitudinal coordinates $\mathbf{x} = (x^1, \dots, x^p)$ and the transverse coordinates $\mathbf{y} = (y^1, \dots, y^{n+1})$, located at $\mathbf{y} = \mathbf{0}$, has the following form:

$$\begin{aligned} G_{MN} dx^M dx^N &= H_p^{-\frac{4(n-1)}{(p+n)\Delta_p}} \left[-dt^2 + d\mathbf{x} \cdot d\mathbf{x} \right] + H_p^{\frac{4(p+1)}{(p+n)\Delta_p}} d\mathbf{y} \cdot d\mathbf{y}, \\ e^{2\phi} &= H_p^{\frac{(p+n)a_p}{\Delta_p}}, \quad F_n = \star(dH_p \wedge dt \wedge dx_1 \wedge \dots \wedge dx_p), \\ H_p &= 1 + \frac{Q_p}{|\mathbf{y}|^{n-1}}, \quad \Delta_p \equiv \frac{(p+n)a_p^2}{2} + \frac{2(p+1)(n-1)}{p+n}. \end{aligned} \quad (2)$$

Such dilatonic p -brane covers all the possible single charged p -branes in string theories. We assume that the p -brane is delocalized in the n -dimensional subspace $\mathcal{K}_n$ with the coordinates y^i ($i = 1, \dots, n$) of the transverse space, where $\mathcal{K}_n$ is a Ricci flat compact manifold with the metric $d\tilde{s}_n^2 = g_{ij}(y^k) dy^i dy^j$. Then, the metric of the delocalized solution has the following form:

$$G_{MN} dx^M dx^N = H_p^{-\frac{4(n-1)}{(p+n)\Delta_p}} \left[-dt^2 + d\mathbf{x} \cdot d\mathbf{x} \right] + H_p^{\frac{4(p+1)}{(p+n)\Delta_p}} \left[d\tilde{y}^2 + g_{ij} dy^i dy^j \right], \quad (3)$$

where $\tilde{y} := y^{n+1}$ and the harmonic function is now given by $H_p = 1 + \tilde{Q}_p |\tilde{y}|$. When $\tilde{Q}_p < 0$, the zeros $\tilde{y} = \pm \tilde{Q}_p^{-1}$ of H_p correspond to the curvature singularities. When $\tilde{Q}_p > 0$, there is no singularity except at $\tilde{y} = 0$, where the delocalized p -brane is located. For the purpose of studying the KK modes of bulk fields, it is convenient to transform the transverse coordinate $\tilde{y}$ to a new one y with which the $(p+2)$ -dimensional part of the metric (with the coordinates (x^μ, y)) takes the standard form of the RS model with the warp factor $\mathcal{W}$. We showed [25] that bulk graviton can be localized on the delocalized p -brane when there are singularities on both sides of the brane. For such case, the metric in the new coordinates is given by

$$\begin{aligned} G_{MN} dx^M dx^N &= \mathcal{W} \left[-dt^2 + d\mathbf{x} \cdot d\mathbf{x} \right] + dy^2 + \mathcal{W}^{-\frac{p+1}{n-1}} g_{ij} dy^i dy^j, \\ \mathcal{W} &= [1 - K_p |y|]^{-\frac{4(n-1)}{2(p+1)+(p+n)\Delta_p}} = [1 - K_p |y|]^{-\frac{8(n-1)}{(p+n)^2 a_p^2 + 4(p+1)n}}, \end{aligned}$$

$$\begin{aligned}
K_p &\equiv -\frac{2(p+1) + (p+n)\Delta_p}{(p+n)\Delta_p} \tilde{Q}_p \\
&= -\frac{(p+n)^2 a_p^2 + 4(p+1)n}{(p+n)^2 a_p^2 + 4(p+1)(n-1)} \tilde{Q}_p.
\end{aligned} \tag{4}$$

Note, the parameter K_p is always positive when $\tilde{Q}_p < 0$.

First, we study the localization properties of bosonic bulk fields. All the bosonic fields can be collectively considered as q -form potentials, since scalar and $U(1)$ gauge fields can be respectively regarded as 0- and 1-form potentials. The action for a massless q -form potential $A_{M_1 \dots M_q}$ with the field strength $F_{M_1 \dots M_{q+1}} = (q+1)\partial_{[M_1} A_{M_2 \dots M_{q+1}]}$ in the D -dimensional bulk spacetime is given by

$$S_q = -\frac{1}{2 \cdot (q+1)!} \int d^D x \sqrt{-G} G^{M_1 N_1} \dots G^{M_{q+1} N_{q+1}} F_{M_1 \dots M_{q+1}} F_{N_1 \dots N_{q+1}}, \tag{5}$$

from which we obtain the following equations of motion for $A_{M_1 \dots M_q}$:

$$\frac{1}{\sqrt{-G}} \partial_{M_1} \left[\sqrt{-G} G^{M_1 N_1} \dots G^{M_{q+1} N_{q+1}} F_{N_1 \dots N_{q+1}} \right] = 0. \tag{6}$$

We take the following KK mode ansatz for the bulk q -form potential:

$$A_{\mu_1 \dots \mu_q}(x^\mu, y, y^i) = a_{\mu_1 \dots \mu_q}^{(m)}(x^\mu) u_m(y), \tag{7}$$

where $a_{\mu_1 \dots \mu_q}^{(m)}$ is assumed to satisfy the following field equations for a massive q -form potential in $(p+1)$ -dimensional flat spacetime:

$$\partial^{\mu_1} f_{\mu_1 \dots \mu_{q+1}}^{(m)} + m^2 a_{\mu_2 \dots \mu_{q+1}}^{(m)} = 0, \tag{8}$$

along with the gauge condition $\partial^{\mu_1} a_{\mu_1 \dots \mu_q}^{(m)} = 0$, where $f_{\mu_1 \dots \mu_{q+1}}^{(m)} \equiv (q+1)\partial_{[\mu_1} a_{\mu_2 \dots \mu_{q+1}]}^{(m)}$ is the field strength of $a_{\mu_1 \dots \mu_q}^{(m)}$. Then, from Eq. (6) we obtain the following Sturm-Liouville equation satisfied by $u_m(y)$:

$$\partial_y \left[\mathcal{W}^{-\frac{p+1}{2(n-1)}-q} \partial_y u_m \right] = m^2 \mathcal{W}^{-\frac{p+1}{2(n-1)}-q-1} u_m, \tag{9}$$

from which we see that the eigenfunctions u_m with different eigenvalues m^2 are orthogonal to one another w.r.t. the weighting function $w(y) = \mathcal{W}^{-\frac{p+1}{2(n-1)}-q-1}$, provided appropriate boundary conditions on u_m and u'_m are satisfied. In the case of the KK zero mode ($m=0$), the general solution to Eq. (9) is given by

$$u_0(y) = c_1 + c_2 \mathcal{W}^{\frac{p+1}{2(n-1)}+q-\frac{(p+n)^2 a_p^2 + 4(p+1)n}{8(n-1)}}, \tag{10}$$

where c_1 and c_2 are integration constants. It can be shown that the y -dependent term in Eq. (10) gives rise to the mass term in the effective action of the q -form potential,

which is contradictory to the fact that u_0 is the KK zero mode. Furthermore, the boundary condition on $\tilde{u}'_0$, where $\tilde{u}_0 \equiv \mathcal{W}^{-\frac{p+1}{4(n-1)} - \frac{q}{2}} u_0$, at $y = 0$, resulting from the δ -function potential in the Schrödinger equation satisfied by $\tilde{u}_0$, requires that $c_2 = 0$. So, the consistent KK zero mode is independent of y . This KK zero mode is normalizable, if its norm

$$\int_{-1/K_p}^{1/K_p} dy u_0^2(y) w(y) \sim (1 - K_p |y|)^{\frac{8(q+1)(n-1) + 4(p+1)(n+1) + (p+n)^2 a_p^2}{(p+n)^2 a_p^2 + 4(p+1)n}} \Big|_{-1/K_p}^{1/K_p} \quad (11)$$

is finite, which is the case when

$$8(q+1)(n-1) + 4(p+1)(n+1) + (p+n)^2 a_p^2 > 0. \quad (12)$$

Note, the $n = 0$ case corresponds to the dilatonic domain wall (with codimension one). For such case, for the KK zero mode of the form potential with the rank $q > (p-1)/2$ to be normalizable, a large enough value of a_p is required [26], as can be seen also from Eq. (12). However, when the codimension of the brane is higher than one ($n \geq 1$), the KK zero mode of bulk form potential of any ranks $q \leq p$ is normalizable for any value of the dilaton coupling parameter a_p . We proposed in Ref. [26] that we have to consider the quantity $u_0^2(y)w(y)$ for determining distribution of the bulk field KK zero mode across the extra spatial direction. With this criterion we see that field distribution of the KK zero mode of the bulk form potential of any ranks is localized around the brane (with the codimension higher than one) for any values of a_p . So, the delocalized p -branes localize bulk form potentials of any ranks in a strict sense, namely, the KK zero mode is normalizable and its field distribution is localized around the brane.

For completeness, we study the dimensional reduction of the bulk form potential to a form potential of lower rank. First, we consider the KK zero mode ansatz of the following form:

$$A_{\mu_1 \dots \mu_r i_1 \dots i_{q-r}}(x^\mu, y, y^i) = a_{\mu_1 \dots \mu_r}(x^\mu) u(y), \quad (13)$$

with all the other components vanishing, where $q - r \leq n$. Following the similar procedure as above, we find that the KK modes are orthogonal w.r.t. the weighting function $w(y) = \mathcal{W}^{\frac{(2q-2r-1)(p+1)}{2(n-1)} - r - 1}$ and the consistent KK zero mode is independent of y . So, the KK zero mode is normalizable, if the following condition is satisfied:

$$8(r+1)(n-1) + 4(p+1)(n+1 - 2q + 2r) + (p+n)^2 a_p^2 > 0. \quad (14)$$

As long as $q - r < (r+1)(n-1)/(p+1) + (n+1)/2$, the KK zero mode is normalizable for any values of a_p . Otherwise, large enough value of a_p^2 is required for normalization of the KK zero mode. By considering the quantity $u_0^2(y)w(y)$, we see that the field distribution for the KK zero mode is localized around the brane, when

$$2(r+1)(n-1) + (p+1)(1 - 2q + 2r) > 0. \quad (15)$$

For such case, the KK zero mode is also normalizable for any values of a_p and therefore the brane localizes the bulk q -form potential in a strict sense. Second, we consider the following form of the KK zero mode ansatz:

$$A_{\mu_1 \dots \mu_r y i_1 \dots i_{q-r-1}}(x^\mu, y, y^i) = a_{\mu_1 \dots \mu_r}(x^\mu) u(y), \quad (16)$$

with all the other components taken to vanish, where $q - r - 1 \leq n$. By substituting this ansatz into Eq. (6), we obtain the following equation satisfied by $u(y)$:

$$\partial_y \left[\mathcal{W}^{\frac{(p+1)(2q-2r-3)}{2(n-1)} - r - 1} u \right] = 0, \quad (17)$$

from which we obtain $u \sim \mathcal{W}^{\frac{(p+1)(3-2q+2r)}{2(n-1)} + r + 1}$. To determine the normalizability, we substitute this KK zero mode ansatz into the bulk action (5). The resulting action has the following form:

$$\begin{aligned} S_q \sim & -\frac{1}{2 \cdot (r+1)!} \int_{-1/K_p}^{1/K_p} dy (1 - K_p |y|)^{\frac{4(2q-2r-3)(p+1) - 8(r+1)(n-1)}{(p+n)^2 a_p^2 + 4(p+1)n}} \\ & \times \int d^{p+1} x \eta^{\alpha_1 \beta_1} \dots \eta^{\alpha_{r+1} \beta_{r+1}} f_{\alpha_1 \dots \alpha_{r+1}} f_{\beta_1 \dots \beta_{r+1}}, \end{aligned} \quad (18)$$

from which we see that the KK zero mode is normalizable when

$$4(n-3+2q-2r)(p+1) - 8(r+1)(n-1) + (p+n)^2 a_p^2 > 0. \quad (19)$$

So, when $q - r - 1 > (n-1)(2r-p+1)/2(p+1)$, the KK zero mode is normalizable for any a_p . Otherwise, large enough a_p^2 is required for the normalization. From the integrand of the y integral in Eq. (18), we see that furthermore the field distribution of the KK zero mode is localized around the brane when $q-r-3/2 > (r+1)(n-1)/(p+1)$, for which case the brane localizes the bulk q -form potential in a strict sense.

Next, we study the localization properties of bulk fermionic fields. First, we consider the massless bulk spin-1/2 fermion Ψ with the action

$$S_{1/2} = \int d^D x \sqrt{-G} i \bar{\Psi} \Gamma^M D_M \Psi, \quad (20)$$

where $D_M \equiv \partial_M + \frac{1}{4} \omega_{MAB} \gamma^{AB}$ is the gravitational covariant derivative on a spinor. Here, ω_{MAB} is the spacetime spin-connection associated with the bulk metric G_{MN} and $\gamma^{AB} \equiv \gamma^{[A} \gamma^{B]}$. The convention for the spacetime vector indices are $M, N, P, \dots$ [$A, B, C, \dots$] for the curved [flat-tangent] bulk spacetime indices, $\mu, \nu, \rho, \dots$ [$\alpha, \beta, \gamma, \dots$] for the curved [flat-tangent] worldvolume spacetime indices, and $i, j, k, \dots$ [$a, b, c, \dots$] for the curved [flat-tangent] transverse space indices. The gamma matrices $\Gamma^M \equiv E_A^M \gamma^A$ with curved indices satisfy $\{\Gamma^M, \Gamma^N\} = 2G^{MN}$, where E_A^M is the inverse of the vielbein E_M^A . From the bulk action (20), we obtain the following equation of motion for the bulk spinor field:

$$\Gamma^M (\partial_M + \frac{1}{4} \omega_{MAB} \gamma^{AB}) \Psi = 0, \quad (21)$$

which reduces to the following form after the bulk metric (3) is substituted:

$$\left[\Gamma^y \left(\partial_y - \frac{p+1}{4(n-1)} \frac{\mathcal{W}'}{\mathcal{W}} \right) + \Gamma^\mu \partial_\mu + \Gamma^m \tilde{D}_m \right] \Psi = 0, \quad (22)$$

where $\tilde{D}_m$ is the gravitational covariant derivative associated with the metric $g_{mn}(y^i)$. In general, the KK mode ansatz for the bulk spinor has the form

$$\Psi(x^\mu, y, y^i) = \psi(x^\mu) u(y) \chi(y^i). \quad (23)$$

Since we are interested in finding the KK zero mode, we impose the zero mode conditions $\gamma^\alpha \partial_\alpha \psi = 0$ and $\Gamma^m \tilde{D}_m \chi = 0$. Then, the bulk equation of motion (22) reduces to the following equation satisfied by $u(y)$:

$$\Gamma^y \left[\partial_y - \frac{p+1}{4(n-1)} \frac{\mathcal{W}'}{\mathcal{W}} \right] u = 0, \quad (24)$$

from which we have $u \sim \mathcal{W}^{\frac{p+1}{4(n-1)}}$. To check the normalizability, we plug this KK zero mode into the bulk action:

$$\begin{aligned} \int d^D x \sqrt{-G} i \bar{\Psi} \Gamma^M D_M \Psi &\sim \int_{-1/K_p}^{1/K_p} dy \mathcal{W}^{-\frac{1}{2}} \int d^{p+1} x i \bar{\psi} \gamma^\alpha \partial_\alpha \psi \\ &= \int_{-1/K_p}^{1/K_p} dy (1 - K_p |y|)^{\frac{4(n-1)}{(p+n)^2 a_p^2 + 4(p+1)n}} \int d^{p+1} x i \bar{\psi} \gamma^\alpha \partial_\alpha \psi. \end{aligned} \quad (25)$$

Note, in the case of the dilatonic domain walls [23], a large enough value of the dilaton coupling parameter is required for localizing the bulk spinor field, as can be seen also from this equation with $n = 0$. On the other hand, in the case of the delocalized p -brane (with $n \geq 1$), the KK zero mode of the bulk fermion is normalizable for any a_p . Furthermore, when $n > 1$, the field distribution of the KK zero mode is localized around the brane as well, so the brane localizes the bulk fermion in a strict sense.

Second, we consider the bulk gravitino Ψ_M with the action

$$S_{3/2} = \int d^D x \sqrt{-G} i \bar{\Psi}_M \Gamma^{MNP} D_N \Psi_P, \quad (26)$$

where $D_M \Psi_N \equiv \partial_M \Psi_N - \Gamma_{MN}^P \Psi_P + \frac{1}{4} \omega_{MAB} \gamma^{AB} \Psi_N$. Here, Γ_{MN}^P is the affine connection associated with G_{MN} and $\Gamma^{MNP} \equiv \Gamma^{[M} \Gamma^N \Gamma^{P]}$. So, the equation of motion for Ψ_M is

$$\Gamma^{MNP} D_N \Psi_P = 0. \quad (27)$$

We choose the gauge $\Psi_y = 0$ and assume $\Psi_m = 0$. In general, the KK mode ansatz then has the form:

$$\Psi_\mu(x^\nu, y, y^i) = \psi_\mu(x^\nu) u(y) \chi(y^i). \quad (28)$$

To obtain the KK zero mode of the bulk gravitino, we impose the zero mode conditions $\gamma^{\alpha\beta\delta}\partial_\beta\psi_\delta = 0$ and $\Gamma^m\tilde{D}_m\chi = 0$, along with the gauge conditions $\partial^\alpha\psi_\alpha = 0$ and $\gamma^\alpha\psi_\alpha = 0$. Then, Eq. (27) reduces to the following equation satisfied by $u(y)$:

$$\Gamma^y \left[\partial_y - \frac{p+2n-1}{4(n-1)} \frac{\mathcal{W}'}{\mathcal{W}} \right] u = 0, \quad (29)$$

from which we have $u \sim \mathcal{W}^{\frac{p+2n-1}{4(n-1)}}$. The bulk action (26) with the KK zero mode substituted takes the form

$$\int d^D x \sqrt{-G} i \bar{\Psi}_M \Gamma^{MNP} D_N \Psi_p \sim \int_{-1/K_p}^{1/K_p} dy \mathcal{W}^{-\frac{1}{2}} \int d^{p+1} x i \bar{\psi}_\alpha \gamma^{\alpha\beta\delta} \partial_\beta \psi_\delta. \quad (30)$$

The integrand of the y -integral has the same form as that in the spinor field effective action (25). So, the normalization and localization properties of the bulk gravitino KK zero mode are the same as those of the bulk spin 1/2 field discussed in the above. Namely, the KK zero mode of the bulk spin 3/2 field is normalizable for any a_p and the delocalized dilatonic p -brane with codimension greater than two ($n > 1$) localizes the bulk spin 3/2 field in a strict sense for any a_p .

References

- [1] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, “The hierarchy problem and new dimensions at a millimeter,” Phys. Lett. **B429** (1998) 263, hep-ph/9803315.
- [2] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, “New dimensions at a millimeter to a Fermi and superstrings at a TeV,” Phys. Lett. **B436** (1998) 257, hep-ph/9804398.
- [3] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension,” Phys. Rev. Lett. **83** (1999) 3370, hep-ph/9905221.
- [4] L. Randall and R. Sundrum, “An alternative to compactification,” Phys. Rev. Lett. **83** (1999) 4690, hep-th/9906064.
- [5] W.D. Goldberger and M.B. Wise, “Bulk fields in the Randall-Sundrum compactification scenario,” Phys. Rev. **D60** (1999) 107505, hep-ph/9907218; “Modulus stabilization with bulk fields,” Phys. Rev. Lett. **83** (1999) 4922, hep-ph/9907447.
- [6] H. Davoudiasl, J.L. Hewett and T.G. Rizzo, “Bulk gauge fields in the Randall-Sundrum model,” Phys. Lett. **B473** (2000) 43, hep-ph/9911262.
- [7] A. Pomarol, “Gauge bosons in a five-dimensional theory with localized gravity,” Phys. Lett. **B486** (2000) 153, hep-ph/9911294.
- [8] B. Bajc and G. Gabadadze, “Localization of matter and cosmological constant on a brane in anti de Sitter space,” Phys. Lett. **B474** (2000) 282, hep-th/9912232.
- [9] Y. Grossman and M. Neubert, “Neutrino masses and mixings in non-factorizable geometry,” Phys. Lett. **B474** (2000) 361, hep-ph/9912408.

- [10] S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, “Bulk standard model in the Randall-Sundrum background,” *Phys. Rev.* **D62** (2000) 084025, hep-ph/9912498.
- [11] I. Oda, “Localization of matters on a string-like defect,” *Phys. Lett.* **B496** (2000) 113, hep-th/0006203; “Bosonic fields in the string-like defect model,” *Phys. Rev.* **D62** (2000) 126009, hep-th/0008012; “Localization of gravitino on a brane,” *Prog. Theor. Phys.* **105** (2001) 667, hep-th/0008134; “Localization of bulk fields on AdS_4 brane in AdS_5 ,” *Phys. Lett. B* **508** (2001) 96, hep-th/0012013; “Locally localized gravity models in higher dimensions,” *Phys. Rev.* **D64** (2001) 026002, hep-th/0102147; “Antisymmetric tensor fields in the locally localized gravity models,” hep-th/0103009; “Kaluza-Klein reduction in a warp geometry revisited,” hep-th/0103158; “Trapping of nonabelian gauge fields on a brane,” hep-th/0103257.
- [12] S. Randjbar-Daemi and M. Shaposhnikov, “Fermion zero-modes on brane-worlds,” *Phys. Lett.* **B492** (2000) 361, hep-th/0008079.
- [13] A. Khoudeir, “A note about localized photons on the brane,” hep-th/0012064.
- [14] M. Tachibana, “Bulk gauge fields in the bigravity model,” hep-th/0101185.
- [15] N. Kaloper, E. Silverstein and L. Susskind, “Gauge symmetry and localized gravity in M theory,” *JHEP* **0105** (2001) 031, hep-th/0006192.
- [16] E. Silverstein, “Gauge fields, scalars, warped geometry, and strings,” *Int. J. Mod. Phys.* **A16** (2001) 641, hep-th/0010144.
- [17] M.J. Duff and J.T. Liu, “Hodge duality on the brane,” hep-th/0010171.
- [18] H. Lu and C.N. Pope, “Branes on the brane,” *Nucl. Phys.* **B598** (2001) 492, hep-th/0008050.
- [19] M.J. Duff, J.T. Liu and W.A. Sabra, “Localization of supergravity on the brane,” hep-th/0009212.
- [20] I. Oda, “A new mechanism for trapping of photon,” hep-th/0103052.
- [21] K. Ghoroku and A. Nakamura, “Massive vector trapping as a gauge boson on a brane,” hep-th/0106145.
- [22] D. Youm, “Solitons in brane worlds,” *Nucl. Phys.* **B576** (2000) 106, hep-th/9911218.
- [23] D. Youm, “Bulk fields in dilatonic and self-tuning flat domain walls,” *Nucl. Phys.* **B589** (2000) 315, hep-th/0002147.
- [24] D. Youm, “Localization of gravity on dilatonic domain walls: Addendum to ‘solitons in brane worlds’,” *Nucl. Phys.* **B596** (2001) 289, hep-th/0007252.
- [25] D. Youm, “Dilatonic p -branes and brane worlds,” *Phys. Rev.* **D62** (2000) 084007, hep-th/0003174.
- [26] D. Youm, “Localization of bulk form fields on dilatonic domain walls,” hep-th/0106240.